

Transit Timing Observations from *Kepler*: VI. Potentially interesting candidate systems from Fourier-based statistical tests

Jason H. Steffen¹, Eric B. Ford², Jason F. Rowe^{8,9}, Daniel C. Fabrycky^{3,4}, Matthew J. Holman⁵, William F. Welsh⁶, Natalie M. Batalha⁷, William J. Borucki⁸, Steve Bryson⁸, Douglas A. Caldwell^{8,9}, David R. Ciardi¹⁰, Jon M. Jenkins^{8,9}, Hans Kjeldsen¹¹, David G. Koch⁸, Andrej Prša¹², Dwight T. Sanderfer⁸, Shawn Seader⁸, Joseph D. Twicken^{8,9},
jsteffen@fnal.gov

ABSTRACT

We analyze the deviations of transit times from a linear ephemeris for the Kepler Objects of Interest (KOI) through Quarter six (Q6) of science data. We conduct two statistical tests for all KOIs and a related statistical test for all pairs of KOIs in multi-transiting systems. These tests identify several systems which show potentially interesting transit timing variations (TTVs). Strong TTV systems have been valuable for the confirmation of planets and their mass measurements. Many of the systems identified in this study should prove fruitful for detailed TTV studies.

Subject headings: planetary systems; planets and satellites: detection, dynamical evolution, and stability; techniques; miscellaneous

1. Introduction

In transiting exoplanet systems, the deviations from a constant orbital period caused by

¹Fermilab Center for Particle Astrophysics, P.O. Box 500, MS 127, Batavia, IL 60510

²Astronomy Department, University of Florida, 211 Bryant Space Sciences Center, Gainesville, FL 32111, USA

³UCO/Lick Observatory, University of California, Santa Cruz, CA 95064, USA

⁴Hubble Fellow

⁵Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁶Astronomy Department, San Diego State University, San Diego, CA 92182-1221, USA

⁷San Jose State University, San Jose, CA 95192, USA

⁸NASA Ames Research Center, Moffett Field, CA, 94035, USA

⁹SETI Institute, Mountain View, CA, 94043, USA

¹⁰NASA Exoplanet Science Institute/California Institute of Technology, Pasadena, CA 91125 USA

¹¹Department of Physics and Astronomy, Aarhus University, DK-8000 Aarhus C, Denmark

¹²Department of Astronomy and Astrophysics, Villanova University, 800 East Lancaster Avenue, Villanova, PA 19085, USA

planet-planet dynamical interactions have proven to be very useful in detecting and characterizing the constituent planets. Transit timing variations (TTVs) for short-timescale and resonant interactions are particularly useful (Agol et al. 2005; Holman and Murray 2005) though secularly induced TTVs have also been studied (Miralda-Escudé 2002; Heyl and Gladman 2007). The *Kepler* mission has profited greatly by the use of TTVs, having used them extensively in the confirmation of a significant fraction of its currently confirmed planets: Kepler-9 (Holman et al. 2010), Kepler-11 (Lissauer et al. 2011), Kepler-18 (Cochran et al. 2011), and Kepler systems 23-34 (Ford et al. 2012a; Fabrycky et al. 2011; Steffen et al. 2012a). TTVs have given: important mass measurements for smaller planets that would be exceedingly difficult to obtain through radial velocity (RV) measurements; stringent limits on the presence of small planets near mean-motion resonance (MMR) with hot Jupiters (Steffen and Agol 2005; Steffen et al. 2012b) with consequent constraints on planet formation and dynamical evolution; and the identifi-

cation of non-transiting planets such as, Kepler-19 (Ballard et al. 2011)—with as-yet undetermined orbital properties.

With the initial release of two quarters of data in 2011 (Borucki et al. 2011) there was an accompanying paper discussing potential TTV candidates (Ford et al. 2011). Now, with the release of more than a year of additional data, we revisit task of identifying systems that show interesting TTV signatures here and in a companion paper (Ford et al. 2012b) using transit times reported in (Rowe et al. 2012). A number of these systems have already been identified, studied, and announced. However, several new systems have been the subject of only cursory investigation and merit further scrutiny. The increased time baseline of the *Kepler* data through Q6 provides many more systems with interesting TTV features as typical timescales for large TTV signals are tens of orbits—which are only now available. With continued operations the scientific yield from *Kepler* and from TTVs specifically will be particularly valuable to the study of planets as many of the orbital and physical properties of planets with small sizes and masses can only be gleaned through TTV analysis. The extension of the *Kepler* mission, with its increased time baseline for TTV studies, will present an opportunity to learn about planets—notably including terrestrial planets and planets near the habitable zone—that will not be rivaled for many decades.

In what follows we present the results of a few straightforward statistical tests on the transit times of known Kepler Objects of Interest (KOI). Two of these tests are applied to all individual KOIs while a third test is applied to pairs of KOIs in the same system in an effort to use correlated TTV signals as a means to improve sensitivity to dynamical interactions. The point of these tests is not to assign rigorous absolute probabilities for the detection of a TTV signal; rather it is to identify systems that merit additional scrutiny. The results of the tests are presented in a single, large table for all analyzed systems (Table 1).

2. Data Reduction

We use KOIs from the planet candidate table reported in (Borucki et al. 2011) and (Batalha et al. 2012). The transit times for each KOI are

determined following the method described in (Ford et al. 2011) and are given in tabular form in (Rowe et al. 2012). Frequently with that method there are some transit times that have anomalously large deviations from a linear ephemeris or that have unusually large error bars. In most cases these anomalies are due to some source of photometric noise or transit detections with low significance rather than from planetary dynamics. However, in cases in which TTVs are larger than the transit duration, the algorithm may fail to find the correct solution, with the optimization algorithm not reaching the global minimum even if the individual transits are highly significant. In those cases, we rerun the algorithm with the initial guesses close enough to the true transit times so the algorithm can model them correctly.

In order to remove non-planetary anomalous TTVs from our analysis, we reduce our timing data by requiring that the mean signal-to-noise ratio (SNR) per transit for a given KOI exceed 3. We eliminate all transit times where the timing residuals deviate from a linear ephemeris by more than four times the Median Absolute Deviation (MAD) of all timing residuals. We also eliminate all transit times where the timing uncertainty is larger than twice the median error of all timing uncertainties for that system. Once this data reduction criteria has been applied, we fit a new linear ephemeris and conduct our remaining analysis on the resulting timing residuals. Table 1 shows the KOIs studied here (1481 with 1009 that satisfy the selection criteria) along with the results of the statistical tests outlined below.

3. Individual Objects

The first analysis we present is applied to each individual KOI. Here we look for a sinusoidal TTV signal with arbitrary amplitude, phase, and period. TTV signals for systems in or near mean-motion resonance (MMR) are expected to be roughly sinusoidal as the description of their motion is dominated by a single sinusoidal component that corresponds to that MMR (Agol et al. 2005). While systems farther from MMR have multi-sinusoidal TTV signatures, the amplitude of the signal is significantly weaker and would not likely be marked as significant in this study (Agol et al. 2005; Steffen 2006). We compare this

single-sinusoid model to the null model where the transit times are fit to a linear ephemeris using the standard F-ratio test with two and five parameters for the null model and the sinusoidal model respectively. This method was first applied to a sample of hot Jupiter candidates in (Steffen et al. 2012b).

We do not claim that the results of this test give an accurate estimate of the statistical significance of a TTV signal, rather that this test can identify interesting systems that merit further scrutiny—as is the goal of this letter. The results of the F-ratio test (p-values that represent the probability that the two-parameter, null model is the correct model of the two models being compared) are shown in Figure 1. We note that the KOIs in light colors in this and all figures are for known planets or well studied KOIs and include: 84 (Kepler-19), 137 (Kepler-18), 157 (Kepler-11), 168 (Kepler-23), 244 (Kepler-25), 250 (Kepler-26), 377 (Kepler-9), 738 (Kepler-29), 806 (Kepler-30), 841 (Kepler-27), 870 (Kepler-28), 935 (Kepler-31), 952 (Kepler-32), and 1102 (Kepler-24).

There are a few planet candidates among the group of well studied objects that have relatively small p-values, notably KOI-806 (Kepler-30) (Tingley et al. 2011; Fabrycky et al. 2011) and KOI-377 (Kepler-9) (Holman et al. 2010). However, it is noteworthy that most of the multi-transiting systems that were studied for anti-correlated TTV signatures in (Ford et al. 2012a; Fabrycky et al. 2011; Steffen et al. 2012a) and the planets in Kepler-11 (Lissauer et al. 2011) do not have particularly low p-values. This is further indication of the importance of multi-transiting planetary systems for the purposes of planet confirmation and dynamical interpretation. A correlation analysis, as is done in (Ford et al. 2012a; Fabrycky et al. 2011; Steffen et al. 2012a) and in the next section of this work, can identify otherwise weak individual candidates. For the F-ratio test, systems that have p-values below 0.02 and that have 10 or more transits are likely to be the most interesting and include (in order of significance) KOIs: 227, 377 (Kepler-9), 806 (Kepler-30), 142, 277 (Kepler-36), 1573, 448, and 918—while still others have undoubtedly interesting signals.

A second analysis for the individual KOIs was to record the ratio of the amplitude of the best

fitting sinusoidal model to the uncertainty in that amplitude (the signal-to-noise ratio (SNR) of the sinusoid amplitude fit). This SNR statistic is given by:

$$SNR = ((A^2 + B^2)(\sigma_A^{-2} + \sigma_B^{-2}))^{1/2} \quad (1)$$

where A and B are the amplitudes of the best fitting sine and cosine components and σ_A and σ_B are the uncertainties in those amplitudes (cf., equation (2) below).

Figure 2 shows the results of this test for all KOIs. As with the F-ratio test, the SNR test shows several systems with sinusoidal amplitudes detected with a large SNR—including many that have not been investigated in depth. The top candidates from this test with SNR greater than 20 and more than 10 transits include (in order of significance) KOIs: 227 (Kepler-36), 377 (Kepler-9), 142, 1573, 806, 984, 13, 277, 473, and 137 (Kepler-18). It is noteworthy that not all of the top candidates from the F-ratio test are top candidates for the SNR test and vice versa. We also point out that in some cases the tests identify some clearly bad candidate systems. For example, the SNR test identifies KOI-13 with high significance—while its timing residuals are caused by an interesting mix of stellar variability and planet-star dynamics, but not an additional planet (Szabó et al. 2011). Figure 3 shows the TTV signal for the three most significant candidates for both the SNR test and the F-ratio test provided they have more than 10 transit times (KOIs 227, 142, and 1573). None of these have yet been claimed as confirmed planets.

4. Object Pairs

For pairs of planets, an additional diagnostic is to look for anticorrelation in the TTV signal. As planets interact with each other we expect many of their TTV signatures to be anticorrelated due to basic effects from conservation of energy and angular momentum—especially when a system is near MMR. It is possible to have short-timescale correlated TTV signals in systems farther from resonance or when there is significant precession in one of the objects (Steffen et al. 2012a; Fabrycky et al. 2011; Ford et al. 2012a). Using an anticorrelation measurement can have much more power to distinguish real TTV signals from noise as one would expect timing noise to be positively correlated




Fig. 1.— Histogram of the p-values resulting from the F-Ratio test. The p-value from known multi-planet systems are the light colored bars. The two most significant systems with this test are KOI-227 and Kepler-9.




Fig. 2.— Histogram of the SNR values resulting from the SNR test. Known multibody systems are the light colored bars. Again, the two most significant systems with this test are KOI-227 and Kepler-9. However, KOI-142 and KOI-1573 also have very large SNR detections.




Fig. 3.— Residuals from a linear ephemeris (observed minus calculated) for the three planet candidates with the most significant results from the SNR test and the F-ratio test that have not yet been claimed as planets and that have more than 10 measured transit times. KOI-227.01 is the top panel, KOI-142.01 is the middle panel, and KOI-1573.01 is the lower panel.

at best and uncorrelated at worst for minimally-interacting planets (some dynamical scenarios can produce positively correlated residuals on short timescales). Following the procedure outlined in (Steffen et al. 2012a) we calculate $\Xi_{\max}$ for each pair of KOIs that satisfy the selection criteria outlined above for all of the systems with multiple transiting objects. This is 155 systems with 262 total pairs that meet the selection criteria. The statistic $\Xi_{\max}$ is found from a periodogram generated by fitting the TTV signal to a set of sinusoidal functions with specific periods:

$$f_i = A \sin\left(\frac{2\pi t}{P_i}\right) + B \cos\left(\frac{2\pi t}{P_i}\right) + C \quad (2)$$

where A , B , and C are model parameters, and P_i is the test timescale¹.

The fitted values for A and B and their measured uncertainties, σ_A and σ_B are then used to calculate Ξ for each of the sampled periods using

$$\Xi = -\left(\frac{A_1 A_2}{\sigma_{A1} \sigma_{A2}} + \frac{B_1 B_2}{\sigma_{B1} \sigma_{B2}}\right) \quad (3)$$

where the “1” and “2” subscripts correspond to the two objects. Finally, the maximum value that Ξ has for a given candidate pair, $\Xi_{\max}$, is the statistic we choose to determine the significance of any interaction between the two objects. The square root of $\Xi_{\max}$ is approximately the significance in terms of σ for Gaussian deviates, though we note that an accurate determination of the significance is best done via Monte Carlo simulation (see Ford et al. 2012a; Fabrycky et al. 2011; Steffen et al. 2012a).

Figure 4 shows the results of this analysis for all KOI pairs. Many of the most significant pairs are in previously identified systems (denoted by the light bars) with Kepler-9 and Kepler-18 being the most significant detections. Still, several have not been previously identified in the literature including KOI-904. Other significant pairs may be found by including systems with fewer transits (e.g., Kepler-30) or with transits detected with less significance, however expanding the analysis down to such systems results in many more spurious

detections. The example of KOI-904 shows how correlation statistics have more power to identify potential TTV systems than simply tests on individual KOIs—the KOI-904 system ranks highly with the correlation test, but the individual objects barely make the top 100 for the F-ratio test and are roughly 50th and 100th for the SNR test. Table 1 includes a subset of all calculated $\Xi_{\max}$ values—specifically, the $\Xi_{\max}$ value found for a given candidate and its neighbor with the next largest orbital period that satisfies the sample selection criteria.




Fig. 4.— Histogram of $\sqrt{\Xi_{\max}}$ for all pairs of candidates. Pairs from well studied multiobject systems are shown with light-colored bars.

The companion paper by Ford et al. (2012b) has similar tests using simple parametric modeling such as quadratic or cubic fits to the TTV signals. Several of the strongest systems using those methods do not appear strong with the Fourier methods and vice-versa—and the relative ranking is quite different. This is to be expected as only in limiting cases do the two models have similar functional form. Moreover, trigonometric functions require a specific relationship between the polynomial powers which generic power-law models do not. Trigonometric functions have a strong physical motivation (as they are the natural basis to solve the equations of motions of gravitating systems) and they allow for more straightforward correlation tests at specific frequencies as is done in this section. Nevertheless, the power-law models of Ford et al. (2012b) can be a faster means of finding interesting systems—especially when there are fewer data to analyze. The full power of the Fourier methods are best realized as multiple cy-

¹There is a typo in (Steffen et al. 2012a) where the period of the planet P is in the denominator of the argument of the sine and cosine functions. This error has been corrected here.

cles exist in the TTV residuals, as will be increasingly the case as more *Kepler* data come available.

5. Conclusions

With the public release of *Kepler* data through Q6, many opportunities are available to study the detailed dynamics of *Kepler* candidate systems. We have conducted some straightforward statistical tests in an effort to identify systems that may be particularly fruitful. Some of the most interesting systems have been identified previously. However, many remain unexplored. Table 1 gives the results of each of these tests and can be used to identify systems that are worthy of additional scrutiny.

Kepler's unique data have enabled unprecedented scientific advances in the study of exoplanets and TTV analyses have played a central role in this endeavor. With these new data and the identification of additional TTV candidate systems several important opportunities are enabled such as the measurement of planetary masses, dynamical studies of mean-motion resonance, the constraints on mutual inclinations, and the discovery of non-transiting exoplanets. These can, in turn, inform models of planet formation and dynamical evolution. Ultimately, TTV analysis is the only means currently available that can measure planet masses of terrestrial planets in the habitable zone of a Sun-like star. At present, its full potential has not been realized.

TABLE 1

^aPERIOD, NUMBER OF TRANSITS, THE RESULTS OF THE SNR OF THE BEST-FIT SINUSOIDAL AMPLITUDE, THE P-VALUE FOR THE F-RATIO TEST, AND $\Xi_{\max}$ FOR EACH KOI.

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
1.01	2.471	149	5.434	0.4032	
2.01	2.205	208	8.227	0.3854	
3.01	4.888	77	6.868	0.3426	
4.01	3.849	84	4.742	0.3937	
5.01	4.78	87	4.3	0.3976	
7.01	3.214	116	4.571	0.4045	
10.01	3.522	127	3.922	0.4126	
12.01	17.855	24	4.152	0.3256	
13.01	1.764	218	28.889	0.1449	
17.01	3.235	142	4.515	0.4091	
18.01	3.548	127	5.887	0.3914	
20.01	4.438	84	5.229	0.3829	
22.01	7.891	57	5.444	0.3555	
41.01	12.816	35	3.725	0.3715	2.711
41.02	6.887	62	4.424	0.3818	6.4
41.03	35.333	12	3.479	0.2422	
42.01	17.834	27	6.33	0.2423	
44.01	66.468	8	3.913	0.1677	
46.01	3.488	132	4.4	0.4074	
49.01	8.314	51	5.031	0.3599	
51.01	10.431	34	4.201	0.3501	
63.01	9.434	48	3.335	0.3922	
64.01	1.951	215	5.474	0.4106	
69.01	4.727	98	4.854	0.3954	
70.01	10.854	40	3.731	0.3782	4.685
70.02	3.696	120	6.514	0.3831	5.301
70.05	19.578	23	5.163	0.2895	
72.01	0.837	432	5.413	0.4211	0.676
72.02	45.294	10	4.663	0.1601	
82.01	16.146	26	4.685	0.3231	2.577
82.02	10.312	40	4.867	0.3538	3.273
82.03	27.453	17	6.311	0.1795	
84.01	9.287	47	10.965	0.197	
85.01	5.86	77	3.937	0.4	1.737
85.02	2.155	185	5.572	0.4056	3.07
85.03	8.131	56	5.517	0.3532	
92.01	65.705	7	8.715	0.0202	
94.01	22.343	9	6.797	0.08	
94.02	10.424	20	7.107	0.2132	2.945
94.04	3.743	50	4.627	0.3626	1.151
97.01	4.885	91	4.002	0.4034	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
98.01	6.79	65	4.241	0.389	
100.01	9.966	44	5.406	0.3331	
102.01	1.735	253	5.612	0.4124	
103.01	14.911	29	18.727	0.0314	
104.01	2.508	166	5.416	0.4048	
105.01	8.981	43	4.571	0.3659	
107.01	7.257	63	3.379	0.4035	
108.01	15.965	26	3.972	0.3443	
110.01	9.941	46	5.351	0.3484	
111.01	11.428	39	4.12	0.3654	2.746
111.02	23.669	19	3.017	0.3551	8.086
111.03	51.757	6	6.253	0.0417	
112.01	51.079	9	7.046	0.0608	
112.02	3.709	115	4.852	0.4018	1.492
115.01	5.412	79	5.253	0.3814	5.958
115.02	7.126	63	4.299	0.3856	
116.01	13.571	33	3.985	0.3644	2.474
116.02	43.845	11	5.062	0.1557	
117.01	14.749	30	4.738	0.3597	
117.02	4.901	88	5.446	0.3826	5.657
117.03	3.18	140	4.693	0.4086	5.176
118.01	24.993	19	4.461	0.2866	
119.01	49.185	10	5.725	0.1164	
122.01	11.523	37	4.098	0.3627	
123.01	6.482	70	4.861	0.3852	2.21
123.02	21.222	22	3.737	0.3293	
124.01	12.691	36	3.749	0.3804	2.512
124.02	31.72	15	4.745	0.2266	
127.01	3.579	122	6.538	0.3807	
128.01	4.943	88	5.743	0.3788	
131.01	5.014	91	4.535	0.3966	
135.01	3.024	150	4.548	0.4099	
137.01	7.642	65	24.769	0.0449	262.344
137.02	14.859	33	20.704	0.0279	
137.03	3.505	120	5.058	0.3999	3.854
138.01	48.938	10	4.115	0.2169	
141.01	2.624	167	4.99	0.4091	
142.01	10.951	41	47.818	0.0017	
144.01	4.176	88	6.022	0.3737	
148.01	4.778	92	4.231	0.4008	16.017
148.02	9.674	48	8.853	0.2421	1.711

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
148.03	42.896	11	4.27	0.2197	
149.01	14.557	29	4.701	0.3382	
150.01	8.409	54	4.807	0.3701	4.551
150.02	28.574	11	5.433	0.1846	
151.01	13.447	30	4.635	0.3238	
152.01	52.091	10	5.213	0.1437	
152.02	27.403	17	6.745	0.181	1.704
152.03	13.484	34	3.9	0.3582	9.329
153.01	8.925	50	5.255	0.3552	
153.02	4.754	93	4.224	0.4026	3.674
155.01	5.661	77	5.168	0.3806	
156.01	8.041	52	4.974	0.361	6.293
156.02	5.189	76	5.063	0.3811	2.211
156.03	11.776	39	4.934	0.342	
157.01	13.025	33	5.389	0.3114	10.359
157.02	22.687	19	4.448	0.2823	4.552
157.03	31.996	13	4.14	0.2282	29.484
157.04	46.69	9	16.842	0.0304	
157.06	10.304	43	5.51	0.3354	8.485
159.01	8.991	46	4.896	0.3504	
161.01	3.106	144	6.301	0.3917	
162.01	14.006	31	3.628	0.3653	
163.01	11.12	42	4.563	0.3575	
165.01	13.222	34	3.231	0.3804	
166.01	12.493	34	5.716	0.3098	
167.01	4.92	90	4.26	0.4008	
168.01	10.742	43	10.497	0.1768	
171.01	5.969	73	5.215	0.3809	1.683
171.02	13.071	34	5.413	0.3256	
172.01	13.722	33	4.215	0.3529	
173.01	10.061	43	3.64	0.3769	
174.01	56.355	8	9.311	0.0298	
176.01	30.23	16	4.712	0.2442	
177.01	21.061	23	3.881	0.3291	
179.01	20.74	21	4.058	0.3253	
180.01	10.046	45	6.378	0.3118	
183.01	2.684	169	4.464	0.413	
186.01	3.243	141	5.741	0.3953	
187.01	30.883	14	4.814	0.2294	
188.01	3.797	118	4.239	0.4087	
189.01	30.36	14	4.92	0.2213	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
190.01	12.265	30	7.197	0.2319	
191.01	15.359	25	3.702	0.347	-0.005
191.02	2.418	147	4.532	0.4098	1.488
191.04	38.651	10	6.899	0.0955	
192.01	10.291	38	3.383	0.3845	
193.01	37.59	13	4.31	0.2306	
194.01	3.121	135	5.25	0.3998	
195.01	3.218	128	3.92	0.4129	
196.01	1.856	243	5.441	0.4127	
197.01	17.276	26	6.53	0.2479	
199.01	3.269	138	4.445	0.4092	
200.01	7.341	57	4.367	0.382	
201.01	4.225	100	4.496	0.4015	
202.01	1.721	254	5.322	0.4145	
203.01	1.486	302	4.166	0.4223	
204.01	3.247	131	5.025	0.4027	
205.01	11.72	39	3.682	0.3829	
206.01	5.334	86	4.788	0.3903	
208.01	3.004	122	4.624	0.406	
209.01	50.791	7	7.197	0.0166	
209.02	18.796	24	4.336	0.3265	0.956
212.01	5.696	48	4.03	0.3818	
214.01	3.312	115	5.248	0.3964	
216.01	20.172	21	4.211	0.2808	
217.01	3.905	117	5.48	0.3934	
219.01	8.025	56	5.05	0.3662	
220.01	2.422	189	5.104	0.4095	
221.01	3.413	131	4.134	0.4115	
222.01	6.312	66	3.888	0.3974	4.62
222.02	12.795	30	3.918	0.3488	
223.01	3.177	136	4.61	0.4061	0.57
223.02	41.007	10	3.358	0.235	
225.01	0.839	335	7.091	0.4099	
226.01	8.309	57	4.485	0.379	
227.01	17.679	21	262.562	7.8356×10^{-7}	
229.01	3.573	126	4.189	0.4095	
232.01	12.466	37	2.861	0.3972	3.915
232.02	5.766	77	4.91	0.3838	1.505
232.03	21.588	20	4.676	0.2658	2.123
232.04	37.997	11	6.816	0.1006	10.749
232.05	56.255	9	3.24	0.2405	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
234.01	9.614	52	4.919	0.3664	
235.01	5.633	81	4.594	0.3919	
237.01	8.508	51	4.508	0.3711	
238.01	17.232	26	3.6	0.3572	
239.01	5.641	74	4.799	0.3869	
240.01	4.287	106	4.495	0.4017	
241.01	13.821	28	6.24	0.26	
242.01	7.258	62	4.568	0.3819	
244.01	12.72	32	4.943	0.3163	
244.02	6.239	69	9.525	0.2574	25.533
245.01	39.792	12	4.023	0.2399	1.861
245.02	21.302	18	4.235	0.2833	6.303
245.04	51.198	6	3.762	0.04	
246.01	5.399	65	4.286	0.3877	
247.01	13.815	24	3.841	0.3205	
248.01	7.204	62	11.535	0.2168	54.409
248.02	10.913	37	9.913	0.1998	
249.01	9.549	41	3.959	0.369	
250.01	12.283	34	6.503	0.285	24.613
250.02	17.251	24	5.075	0.2774	1.428
250.04	46.828	9	3.469	0.2354	
251.01	4.164	100	4.249	0.4038	
252.01	17.604	24	3.791	0.3284	
253.01	6.383	57	4.462	0.3797	
254.01	2.455	174	7.189	0.3885	
255.01	27.522	17	3.309	0.3347	
256.01	1.379	233	6.551	0.4045	
257.01	6.883	51	4.583	0.3655	
258.01	4.158	88	6.239	0.371	
260.01	10.496	36	5.289	0.3314	
261.01	16.238	27	6.192	0.2619	
262.01	7.813	56	4.01	0.388	6.515
262.02	9.376	45	4.141	0.3727	
263.01	20.719	18	3.107	0.3453	
269.01	18.011	22	6.069	0.2336	
270.01	12.583	34	3.954	0.3546	3.182
270.02	33.673	12	4.472	0.2303	
271.01	48.631	8	4.211	0.1331	
271.02	29.392	16	4.507	0.2536	6.753
273.01	10.574	38	6.158	0.2923	
275.01	15.792	20	3.442	0.3389	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
276.01	41.746	11	3.787	0.2364	
277.01	16.228	29	27.532	0.0079	
279.01	28.455	15	6.483	0.153	
279.02	15.413	24	4.031	0.3267	2.809
280.01	11.873	36	3.063	0.3887	
281.01	19.557	19	7.547	0.2415	
282.01	27.508	14	4.884	0.2091	
283.01	16.092	27	4.091	0.3361	
284.01	18.01	19	4.484	0.2789	
284.02	6.415	49	4.411	0.3665	2.715
284.03	6.178	54	3.897	0.3896	4.607
285.01	13.749	33	4.761	0.3392	
288.01	10.275	37	5.953	0.3091	
289.01	26.629	13	4.156	0.2629	
291.01	31.518	15	4.542	0.2469	
292.01	2.587	143	5.061	0.4013	
294.01	34.436	11	3.895	0.2332	
295.01	5.317	68	5.766	0.3595	
296.01	28.862	11	3.434	0.2541	
297.01	5.652	67	4.452	0.381	
298.01	19.964	22	3.251	0.3562	
299.01	1.542	256	4.621	0.4184	
301.01	6.003	73	5.231	0.3797	
302.01	24.855	11	3.773	0.2585	
303.01	60.929	6	5.541	0.0464	
304.01	8.512	48	3.838	0.382	
305.01	4.604	90	4.242	0.3998	
306.01	24.308	15	5.159	0.2297	
307.01	19.674	25	3.928	0.3339	
308.01	35.595	12	13.358	0.0315	
312.01	11.579	37	7.49	0.2575	3.893
312.02	16.399	27	4.914	0.2998	
313.01	18.736	20	4.091	0.3135	
313.02	8.436	51	5.866	0.3329	8.154
314.01	13.781	31	5.212	0.3352	4.86
314.02	23.09	19	3.973	0.314	
315.01	35.591	11	3.851	0.2505	
317.01	22.208	19	3.956	0.3029	
318.01	38.584	11	4.927	0.1716	
319.01	46.151	10	9.571	0.0327	
321.01	2.426	180	4.997	0.4105	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
323.01	5.836	71	5.593	0.3681	
330.01	7.974	54	3.539	0.3936	
331.01	18.684	24	4.004	0.3381	
332.01	5.459	81	5.742	0.3749	
333.01	13.285	31	4.198	0.3337	
335.01	46.567	9	6.244	0.0859	
337.01	19.784	24	4.252	0.3185	
338.01	7.011	57	4.417	0.3776	
339.01	1.98	196	4.328	0.4159	1.721
339.02	12.834	34	5.345	0.3146	
340.01	23.673	16	4.856	0.2379	
341.01	7.171	58	3.386	0.3992	
343.01	4.762	90	4.576	0.3977	4.043
343.02	2.024	204	5.826	0.406	6.136
343.03	41.81	10	4.837	0.1532	
344.01	39.309	12	4.598	0.1963	
345.01	29.886	12	4.684	0.2125	
346.01	12.925	34	3.491	0.3794	
348.01	28.511	16	5.986	0.1743	
349.01	14.387	28	4.677	0.3153	
350.01	12.991	28	4.013	0.344	
351.03	59.74	7	7.891	0.0131	
352.01	27.083	17	4.073	0.2749	
354.01	15.96	28	4.331	0.3358	
355.01	4.903	88	4.12	0.4017	
356.01	1.827	237	5.891	0.409	
361.01	3.248	130	6.291	0.3877	
366.01	75.112	6	3.314	0.1796	
367.01	31.579	11	5.807	0.1375	
370.01	42.882	10	4.391	0.1929	
370.02	22.951	18	5.78	0.2145	4.017
377.01	19.265	24	95.814	0.0006	1755.65
377.02	38.882	13	70.098	0.0001	
379.01	6.717	66	4.18	0.3948	
384.01	5.08	65	4.853	0.393	
385.01	13.145	34	4.592	0.3427	
386.01	31.158	11	6.237	0.2416	
387.01	13.9	30	4.461	0.3372	
388.01	6.15	65	4.482	0.3878	
392.01	33.418	12	7.155	0.0949	
393.01	21.416	17	3.913	0.298	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
398.01	51.847	8	4.336	0.1388	
398.02	4.18	101	4.742	0.3987	3.824
401.01	29.199	14	6.666	0.1292	
403.01	21.057	19	4.165	0.2991	
408.01	7.382	58	4.001	0.3927	4.004
408.02	12.561	32	3.698	0.3719	1.627
408.03	30.826	14	5.204	0.2104	
409.01	13.249	31	4.149	0.3496	
410.01	7.217	58	3.797	0.3937	
412.01	4.147	89	4.165	0.4007	
413.01	15.229	29	4.899	0.3073	2.113
413.02	24.675	18	3.949	0.2991	
416.01	18.208	20	3.883	0.3158	3.891
416.02	88.254	6	5.653	0.0549	
417.01	19.193	22	4.378	0.3138	
418.01	22.418	16	4.97	0.2204	
419.01	20.132	21	4.436	0.293	
420.01	6.01	74	4.397	0.3935	
421.01	4.454	101	4.893	0.397	
423.01	21.087	18	4.438	0.2958	
425.01	5.428	71	5.676	0.376	
426.01	16.301	26	4.048	0.3313	
427.01	24.615	16	6.981	0.2122	2.077
427.02	42.951	12	4.15	0.2441	
428.01	6.873	65	3.672	0.4	
429.01	8.6	55	3.271	0.3999	
430.01	12.377	27	2.989	0.3737	
431.01	18.87	22	4.006	0.3051	1.098
431.02	46.903	10	4.035	0.2238	
432.01	5.263	83	4.011	0.4019	
435.01	20.55	18	4.508	0.2888	
438.01	5.931	73	4.999	0.3805	2.065
438.02	52.663	8	4.934	0.099	
439.01	1.902	229	4.831	0.4156	
440.01	15.907	28	4.856	0.3106	
440.02	4.973	85	5.297	0.3843	3.807
442.01	13.54	33	4.173	0.356	
443.01	16.218	27	5.971	0.2691	
444.01	11.723	33	4.306	0.3485	
446.01	16.709	22	4.742	0.298	6.528
446.02	28.551	12	3.864	0.2496	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
448.01	10.14	43	5.773	0.3207	11.975
448.02	43.611	10	17.921	0.0121	
452.01	3.706	121	3.573	0.4145	
454.01	29.007	11	5.388	0.1651	
456.01	13.7	30	19.024	0.2226	
457.01	4.921	87	4.293	0.3977	3.419
457.02	7.064	65	7.217	0.3198	
458.01	53.718	6	11.63	0.0059	
459.01	19.446	19	6.004	0.242	
460.01	17.587	23	5.958	0.3361	
463.01	18.478	22	5.395	0.2701	
464.01	58.362	8	7.338	0.0408	
464.02	5.35	79	4.424	0.3941	1.95
466.01	9.391	50	5.029	0.3563	
467.01	18.009	23	3.932	0.3355	
468.01	22.184	19	4.69	0.3059	
469.01	10.329	41	4.424	0.3654	
470.01	3.751	113	4.89	0.3995	
471.01	21.348	21	4.183	0.3139	
472.01	4.244	102	4.849	0.3947	
473.01	12.706	35	24.84	0.1392	
474.01	10.946	39	3.231	0.3882	3.203
474.02	28.986	14	4.671	0.2434	
475.01	8.181	54	4.33	0.38	3.86
475.02	15.313	29	3.566	0.3694	
476.01	18.428	24	3.861	0.2986	
477.01	16.543	25	6.145	0.2378	
478.01	11.023	39	3.612	0.3826	
479.01	34.189	12	5.052	0.1697	
480.01	4.302	108	4.231	0.4068	
481.01	7.65	58	4.468	0.3805	2.665
481.03	34.261	12	3.5	0.2634	
483.01	4.799	72	3.612	0.4028	
484.01	17.205	26	5.619	0.2682	
486.01	22.183	20	5.768	0.2384	
487.01	7.659	59	4.826	0.3775	
488.01	9.379	46	4.699	0.3572	
490.03	7.406	59	3.967	0.3926	
492.01	29.911	12	4.989	0.1917	
494.01	25.697	15	7.492	0.2474	
496.01	1.617	253	5.056	0.4159	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
497.01	13.193	27	4.634	0.3185	29.521
499.01	9.668	36	4.515	0.3523	
500.01	7.053	49	7.014	0.3013	
500.02	9.522	38	8.892	0.2127	
501.01	24.795	17	3.688	0.3152	
503.01	8.222	45	3.788	0.3786	
504.01	40.606	6	7.74	0.0291	
505.01	13.767	27	6.223	0.2491	2.577
506.01	1.583	269	5.683	0.4135	
507.01	18.492	21	4.219	0.304	
508.01	7.931	43	3.952	0.3797	4.24
508.02	16.666	19	4.28	0.3044	
509.01	4.167	103	4.348	0.4037	4.536
509.02	11.464	38	3.683	0.3663	
510.01	2.94	120	4.744	0.4006	2.448
510.02	6.389	49	3.123	0.4005	
510.03	14.627	22	4.995	0.2817	
511.01	8.006	54	4.429	0.3788	4.666
512.01	6.51	66	4.557	0.3815	
513.01	35.181	14	3.025	0.3407	
517.01	2.752	150	4.759	0.4077	3.955
518.01	13.982	27	3.56	0.3542	
518.02	44	10	3.508	0.2238	
519.01	11.903	35	4.139	0.3622	5.972
519.02	34.036	14	4.292	0.2213	
520.01	12.76	31	3.662	0.3653	
520.03	25.751	18	4.168	0.2913	4.757
521.01	10.161	44	3.257	0.3942	
522.01	12.83	33	5.912	0.2955	
523.01	49.413	9	4.323	0.1843	1.571
523.02	36.854	12	4.954	0.1877	
524.01	4.592	96	18.435	0.1451	
525.01	11.532	38	4.748	0.3421	0.388
526.01	2.105	201	4.852	0.4127	
528.01	9.577	48	6.222	0.3253	
528.02	96.672	6	10.154	0.0042	0.388
528.03	20.553	22	4.287	0.3056	
530.01	10.94	37	4.147	0.3606	
531.01	3.687	100	3.552	0.4115	0.2911
532.01	4.222	106	4.323	0.4028	
533.01	16.55	27	5.241	0.2911	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
534.01	6.4	64	3.759	0.3966	
535.01	5.853	75	4.555	0.3897	
537.01	2.82	135	5.046	0.4036	
538.01	21.217	19	3.645	0.3286	
541.01	13.646	32	4.07	0.3467	
542.01	41.886	11	3.768	0.2443	
543.01	4.302	98	3.485	0.4105	
546.01	20.684	19	3.687	0.3212	
547.01	25.303	19	4.359	0.2881	
548.01	21.3	19	4.12	0.3098	
550.01	13.024	25	3.267	0.3627	
551.01	11.637	32	5.204	0.3125	
552.01	3.055	130	5.368	0.399	
554.01	3.658	121	5.126	0.3989	
557.01	15.656	24	3.278	0.3649	
558.01	9.178	43	5.444	0.3391	
560.01	23.675	17	4.34	0.2845	
561.01	5.379	81	4.639	0.3892	
563.01	15.284	30	7.088	0.2294	
564.01	21.055	19	4.625	0.2782	
566.01	25.855	9	5.001	0.1607	
567.01	10.688	37	4.802	0.3383	5.153
567.02	20.303	23	5.34	0.2817	1.285
567.03	29.023	14	3.715	0.2881	
569.01	20.729	20	7.87	0.1586	
571.01	7.267	56	6.161	0.3433	3.026
571.02	13.343	32	3.1	0.3594	2.437
571.04	22.408	20	3.821	0.2801	
572.01	10.64	41	4.136	0.367	
573.01	5.997	75	4.182	0.3938	
574.01	20.135	23	3.662	0.3491	
575.01	24.316	16	4.787	0.2681	
578.01	6.412	69	3.667	0.4022	
580.01	6.521	66	4.759	0.3826	
581.01	6.997	60	5.155	0.3668	
582.01	5.945	73	4.794	0.3872	4.649
582.02	17.739	23	4.776	0.2972	
584.01	9.927	43	4.355	0.3686	2.981
584.02	21.223	20	6.116	0.2944	
585.01	3.722	114	6.379	0.3819	
586.01	15.78	22	3.906	0.3449	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
587.01	14.035	32	3.776	0.3575	6.259
588.01	10.356	41	3.872	0.3731	
590.01	11.389	35	3.781	0.3732	
590.02	50.697	8	5.947	0.084	
592.01	39.749	12	8.804	0.0602	
593.01	9.997	43	3.856	0.381	8.225
597.01	17.308	22	3.188	0.3659	
598.01	8.308	56	4.655	0.3715	
599.01	6.454	67	5.399	0.3697	
600.01	3.596	106	4.409	0.4045	
601.01	5.404	76	3.934	0.3979	2.592
601.02	11.679	37	6.289	0.3089	
602.01	12.914	32	3.884	0.3596	
605.01	2.628	163	5.092	0.4063	
607.01	5.894	44	6.674	0.34	
609.01	4.397	91	4.987	0.3913	4.466
610.01	14.283	22	4.16	0.3237	
611.01	3.252	126	4.652	0.4055	
612.01	20.74	16	4.428	0.2775	
612.02	47.426	7	1.624	0.0523	
614.01	12.875	32	5.813	0.3008	3.224
617.01	37.865	10	6.04	0.1363	
618.01	9.071	48	4.386	0.3701	
620.01	45.155	9	5.447	0.1377	
623.01	10.35	42	4.591	0.3576	
623.02	15.677	25	4.671	0.3118	0.266
624.01	17.79	23	3.998	0.3379	
624.02	49.567	9	7.379	0.0604	
625.01	38.139	10	3.473	0.2629	
626.01	14.587	25	4.717	0.3113	
627.01	7.752	57	3.821	0.3953	0.273
628.01	14.486	32	4.516	0.3373	
629.01	40.699	10	3.124	0.2853	
632.01	7.239	57	4.602	0.3757	
635.01	16.72	20	4.117	0.3103	
638.01	23.636	15	2.963	0.266	0.26
639.01	17.98	20	5.272	0.273	
640.01	30.997	13	4.524	0.26	
641.01	14.852	26	4.526	0.3343	
644.01	45.978	7	6.787	0.0512	
645.02	23.784	16	3.733	0.3109	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
647.01	5.169	87	3.856	0.4058	
649.01	23.45	19	4.677	0.2854	
650.01	11.955	38	4.525	0.3575	
652.01	16.081	26	5.036	0.2958	
654.01	8.594	41	4.981	0.3482	
655.01	25.672	17	3.199	0.3251	
657.01	4.069	83	6.761	0.3554	5.688
657.02	16.283	22	4.433	0.3112	
658.01	3.163	133	5.434	0.3987	5.205
658.02	5.371	77	4.849	0.3834	
659.01	23.206	19	4.821	0.2775	
660.01	6.08	60	4.73	0.3744	
661.01	14.401	25	4.018	0.3434	
662.01	10.214	43	4.588	0.361	
663.01	2.756	118	5.008	0.4007	2.645
663.02	20.307	18	4.672	0.2592	
664.01	13.137	34	4.82	0.3322	
665.01	5.868	74	4.62	0.3887	
666.01	22.248	18	4.823	0.2959	
667.01	4.305	97	4.748	0.3975	
670.01	9.489	45	4.8	0.3518	
672.01	16.088	26	4.137	0.3509	0.858
672.02	41.749	10	4.832	0.152	
673.01	4.417	97	3.921	0.4065	
674.01	16.339	15	3.466	0.3063	
676.01	7.973	55	6.682	0.3159	
676.02	2.453	173	3.766	0.418	2.099
678.01	6.04	68	4.168	0.3944	
678.02	4.139	93	4.825	0.3931	3.463
679.01	31.806	12	4.685	0.2216	
680.01	8.6	48	4.365	0.3686	
684.01	4.035	104	3.245	0.4149	
685.01	3.174	134	4.728	0.4058	
686.01	52.514	8	5.964	0.0952	
688.01	3.276	135	3.491	0.4179	
689.01	15.874	15	4.043	0.1792	
691.01	29.666	14	4.299	0.2464	
693.01	28.779	13	5.928	0.1597	
693.02	15.66	27	4.739	0.3143	5.857
694.01	17.421	23	3.234	0.3661	
695.01	29.908	13	4.641	0.2468	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
697.01	3.032	81	5.917	0.3721	
698.01	12.719	32	5.434	0.3132	
700.01	30.864	15	3.305	0.3225	
700.02	9.361	45	7.036	0.2967	2.241
700.03	14.667	28	2.939	0.3852	3.305
701.01	18.164	22	3.88	0.3391	
701.02	5.715	75	4.874	0.3839	2.848
704.01	18.396	20	11.217	0.3273	
707.01	21.776	19	4.459	0.2886	8.616
707.02	41.027	12	4.387	0.2234	
707.03	31.785	14	3.977	0.274	1.423
707.04	13.176	34	4.006	0.3676	2.471
708.01	17.406	23	3.263	0.3535	
708.02	7.693	54	4.78	0.3691	2.776
709.01	21.384	22	4.89	0.3088	
710.01	5.375	77	4.213	0.3947	
711.01	44.699	9	4.135	0.2101	
714.01	4.182	108	3.974	0.4073	
716.01	26.893	16	3.502	0.3174	
717.01	14.707	29	5.545	0.2927	
718.01	4.585	99	4.244	0.4029	4.306
718.02	22.715	18	4.185	0.3086	3.011
718.03	47.906	10	5.317	0.1197	
719.01	9.034	48	4.405	0.3737	
720.01	5.691	68	3.836	0.3982	7.732
720.02	10.042	43	4.834	0.3526	3.849
720.03	18.37	23	5.289	0.2723	
720.04	2.796	147	5.092	0.4022	5.038
721.01	13.724	31	5.298	0.3142	
722.01	46.408	10	3.215	0.2672	
723.01	3.937	102	4.841	0.398	3.294
723.02	28.082	13	7.868	0.0853	
723.03	10.089	39	4.601	0.3516	4.523
725.01	7.305	61	5.843	0.3493	
728.01	7.189	59	5.765	0.3537	
730.01	14.788	26	3.981	0.3432	5.444
730.03	19.722	20	4.098	0.3121	
732.01	1.26	342	4.373	0.4225	
733.01	5.925	72	4.869	0.3831	1.915
733.02	11.349	37	4.131	0.3634	
734.01	24.543	16	5.682	0.1727	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
735.01	22.341	19	5.57	0.2977	26.102
736.01	18.794	22	7.622	0.3124	
737.01	14.498	29	10.639	0.3277	
738.01	10.338	45	10.787	0.2776	
738.02	13.291	32	4.833	0.3388	
740.01	17.672	24	3.938	0.3208	
741.01	23.355	17	4.782	0.2526	
743.01	19.404	22	3.824	0.3319	
745.01	16.47	26	4.2	0.3382	
746.01	9.274	42	3.027	0.3965	
747.01	6.029	69	4.468	0.3919	2.181
749.01	5.35	80	3.793	0.4034	
750.01	21.677	20	6.737	0.1621	
752.01	9.488	46	3.307	0.3935	
752.02	54.415	9	4.836	0.1284	
753.01	19.899	22	5.215	0.2835	
756.01	11.094	37	3.721	0.3722	
757.01	16.069	24	4.591	0.3024	
757.02	41.192	10	4.162	0.1793	
757.03	6.253	59	5.513	0.3592	6.709
758.01	16.013	22	4.486	0.3069	
759.01	32.624	11	4.891	0.1692	
760.01	4.959	86	4.004	0.402	
763.01	19.651	18	5.953	0.2147	
764.01	41.439	11	9.072	0.048	
765.01	8.354	46	4.542	0.3358	
766.01	4.126	99	4.437	0.4022	
767.01	2.817	156	4.211	0.4147	
769.01	4.281	100	4.068	0.406	10.104
773.01	38.378	8	4.075	0.1744	
774.01	7.443	62	6.094	0.3483	
775.01	16.385	23	5.291	0.2592	
775.02	7.877	44	7.56	0.2838	
775.03	36.446	11	7.836	0.1461	
776.01	3.729	122	5.037	0.3992	
777.01	40.42	11	6.548	0.1032	
779.01	10.406	32	4.139	0.3547	
780.01	2.337	174	6.287	0.399	
781.01	11.598	40	5.07	0.345	12.825
782.01	6.575	63	6.056	0.3506	
783.01	7.275	62	4.697	0.3785	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
784.01	19.271	20	16.52	0.059	4.005
785.01	12.393	32	5.051	0.3281	
787.01	4.431	97	4.83	0.3946	
787.02	11.379	39	4.036	0.3676	
788.01	26.394	17	2.686	0.3523	
790.01	8.472	45	3.439	0.3936	
791.01	12.612	35	2.944	0.3911	
795.01	6.77	63	5.101	0.3724	
797.01	10.182	42	3.81	0.3802	
799.01	1.627	222	5.982	0.4061	
800.01	2.711	159	4.314	0.4118	2.096
800.02	7.212	53	3.466	0.3961	
801.01	1.626	219	4.885	0.4151	
802.01	19.62	17	4.214	0.2971	
804.01	9.029	46	5.004	0.3575	
805.01	10.328	32	4.116	0.3564	
806.02	60.325	6	8.23	0.0095	
806.03	29.287	15	39.307	0.0008	153.317
809.01	1.595	201	5.862	0.4053	
810.01	4.783	92	4.658	0.3967	
811.01	20.506	22	4.79	0.2988	
812.01	3.34	127	4.818	0.4041	
812.02	20.061	23	3.888	0.3447	3.818
812.03	46.183	8	6.12	0.0684	
813.01	3.896	93	6.302	0.3656	
814.01	22.366	16	2.781	0.2695	
815.01	34.844	14	7.125	0.1275	
816.01	7.748	38	5.887	0.3051	
817.01	23.968	13	9.828	0.0572	
818.01	8.114	49	4.207	0.3808	
821.01	21.813	13	4.736	0.2155	
822.01	7.919	45	4.71	0.3646	2.835
823.01	1.028	329	10.252	0.3884	
824.01	15.376	6	10.833	0.0046	
825.01	8.104	38	5.804	0.3095	
826.01	6.366	73	3.667	0.4009	
827.01	5.976	62	3.741	0.3923	
829.01	18.648	23	5.7	0.2514	
829.03	38.561	12	3.293	0.2924	
830.01	3.526	126	4.889	0.4016	
833.01	3.951	88	4.135	0.404	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
834.01	23.654	19	6.106	0.2211	
834.02	13.234	31	4.874	0.3223	1.509
835.01	11.763	29	5.907	0.2661	8.451
835.02	56.229	8	4.448	0.1496	
837.01	7.954	48	3.967	0.3823	
838.01	4.859	93	4.266	0.3984	
840.01	3.04	144	5.697	0.3979	
841.01	15.335	29	8.641	0.1753	19.262
841.02	31.33	11	5.067	0.1763	
842.01	12.718	30	3.973	0.3607	0.696
842.02	36.065	12	4.813	0.2082	
843.01	4.19	107	5.471	0.3873	
844.01	3.71	93	4.413	0.4	
845.01	16.33	22	4.345	0.3075	
846.01	27.808	16	6.215	0.1665	
847.01	80.871	6	4.885	0.0668	
849.01	10.355	45	5.258	0.3417	
850.01	10.526	45	3.61	0.3873	
851.01	4.584	98	4.081	0.4055	
853.01	8.204	53	3.641	0.392	10.23
853.02	14.497	30	7.05	0.2808	
854.01	56.056	8	6.722	0.0487	
855.01	41.408	10	5.953	0.1243	
856.01	39.749	11	3.607	0.2669	
857.01	5.715	61	4.209	0.388	
858.01	13.61	28	3.621	0.354	
864.01	4.312	91	4.269	0.4029	5.52
864.02	20.05	22	4.167	0.3154	
867.01	16.085	28	4.992	0.3026	
869.01	7.49	59	4.529	0.3823	3.201
869.02	36.285	12	3.181	0.223	
869.03	17.461	24	5.69	0.2667	0.978
870.01	5.912	72	10.823	0.2513	29.827
870.02	8.986	49	5.827	0.3421	
871.01	12.941	30	4.545	0.3397	
872.01	33.603	14	12.047	0.0313	
874.01	4.602	93	4.701	0.392	
875.01	4.221	102	3.977	0.4085	
876.01	6.998	56	3.714	0.3916	
877.01	5.955	70	3.551	0.4017	4.235
877.02	12.04	35	5.336	0.3169	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
878.01	23.59	17	4.062	0.2899	
880.01	26.442	16	4.722	0.2581	1.727
880.02	51.529	9	4.583	0.1748	
880.03	5.902	75	3.614	0.4038	5.344
881.01	21.023	19	2.842	0.361	
882.01	1.957	225	8.348	0.3828	
883.01	2.689	158	5.226	0.4033	
884.01	9.439	45	4.717	0.3587	4.222
884.02	20.473	18	9.076	0.0853	
886.01	8.012	56	12.379	0.246	
887.01	7.411	59	4.882	0.3659	
889.01	8.885	51	5.279	0.354	
890.01	8.099	56	3.843	0.3911	
891.01	10.007	43	4.104	0.3743	
892.01	10.372	38	3.328	0.3875	
893.01	4.408	77	4.039	0.4019	
895.01	4.409	102	6.172	0.3764	
896.01	16.24	25	5.135	0.2931	
896.02	6.308	69	3.986	0.3978	2.163
897.01	2.052	206	6.095	0.4043	
898.01	9.771	44	5.533	0.3318	7.133
898.02	5.17	83	5.88	0.3715	9.014
898.03	20.09	22	3.83	0.32	
899.01	7.114	60	5.462	0.3544	7.366
899.03	15.368	31	4.425	0.3338	
900.01	13.81	25	3.451	0.3599	
901.01	12.733	34	4.129	0.3671	
903.01	5.007	85	4.662	0.3905	
904.02	27.945	15	7.083	0.1422	30.626
904.03	42.157	10	9.098	0.1469	
905.01	5.795	75	3.586	0.4033	
906.01	7.157	59	3.877	0.3923	3.376
906.02	17.648	23	4.21	0.3204	
907.01	16.514	20	4.509	0.2899	3.206
907.02	30.133	11	5.368	0.1604	
908.01	4.708	95	5.033	0.3896	
910.01	5.392	77	4.618	0.3911	
912.01	10.848	38	4.554	0.3542	
913.01	4.082	107	4.875	0.4012	
916.01	3.315	125	5.408	0.3961	
917.01	6.72	65	4.416	0.3812	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$	
918.01	39.644	10	11.666	0.0186	3.989	
920.01	21.804	19	4.231	0.301		
921.01	10.282	40	5.651	0.321		
921.02	18.119	22	3.86	0.3343		
922.01	5.154	82	4.003	0.4009		
923.01	5.743	78	4.051	0.4009		
924.01	39.476	12	4.861	0.192		
926.01	3.166	122	4.431	0.4051		
929.01	6.492	55	4.966	0.3662		
931.01	3.856	117	4.408	0.4069		
934.01	5.827	77	4.434	0.3921	1.908	
934.03	18.747	20	3.34	0.3498	16.225	
935.01	20.861	23	7.642	0.1883		
935.02	42.633	9	5.199	0.1514		
936.01	9.468	46	4.248	0.3758		
937.01	20.835	18	6.369	0.1998		
938.01	9.946	43	4.39	0.3587		
940.01	6.105	75	5.239	0.3776		
941.01	6.582	70	3.852	0.3997		4.005
941.03	24.665	19	4.453	0.2833		
942.01	11.515	37	3.922	0.3704		
943.01	3.601	121	4.316	0.4055	5.259	
944.01	3.108	147	5.242	0.4043		
945.01	25.847	15	4.27	0.2943		
945.02	40.718	10	6.671	0.1056		
947.01	28.599	14	2.722	0.336		
949.01	12.533	31	4.489	0.3554		
951.01	13.197	35	3.398	0.3804		1.417
951.02	33.653	12	4.239	0.2631		
952.01	5.901	53	6.798	0.3203		28.179
952.02	8.752	41	8.306	0.2758		7.633
952.03	22.781	14	3.532	0.2878	1.848	
953.01	3.584	117	4.086	0.4098		
954.01	8.115	51	4.315	0.376		
954.02	36.926	10	3.942	0.2351		
955.01	7.039	55	4.073	0.3847		
956.01	8.361	52	3.914	0.385		
960.01	15.801	25	6.668	0.244		
961.01	1.214	336	4.783	0.4207		
972.01	13.119	35	4.512	0.3461		
974.01	53.507	9	3.09	0.2403		

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
975.01	2.786	148	4.643	0.4087	
976.01	52.569	9	6.608	0.2361	
977.01	1.354	298	6.448	0.4116	
984.01	4.288	103	33.905	0.0325	
986.01	8.188	54	3.806	0.3884	3.806
986.02	76.051	6	9.15	0.0079	
987.01	3.179	127	3.89	0.4122	
988.01	10.381	40	3.491	0.3844	2.454
988.02	24.57	18	3.458	0.3285	
991.01	12.062	34	3.604	0.3804	
993.01	21.853	21	4.709	0.2912	
999.01	16.568	27	5.125	0.3164	
1001.01	40.811	11	4.576	0.1958	
1003.01	8.361	49	10.766	0.2186	
1005.01	35.618	13	3.289	0.3114	
1014.01	17.317	21	3.994	0.3121	
1015.01	9.429	22	3.839	0.3257	
1017.01	17.445	24	4.224	0.325	
1020.01	54.356	8	6.147	0.0762	
1022.01	18.827	20	3.265	0.3408	
1024.01	5.748	73	4.718	0.3886	
1029.01	32.309	14	3.594	0.2859	
1052.01	17.028	27	4.33	0.3266	
1060.01	12.11	34	4.931	0.3272	
1061.01	41.815	10	3.215	0.2449	
1066.01	5.715	44	4.32	0.3685	
1074.01	3.771	65	3.989	0.3943	
1078.01	3.354	129	5.877	0.392	7.842
1078.02	6.877	66	4.389	0.388	6.362
1078.03	28.463	15	4.448	0.2583	
1081.01	9.956	47	14.976	0.2075	
1086.01	27.666	14	4.912	0.2365	
1089.02	12.218	31	4.003	0.352	
1094.01	6.1	69	4.694	0.3803	
1095.01	51.598	8	7.2	0.0275	
1102.01	12.333	37	8.927	0.2043	
1102.02	8.145	51	8.892	0.2587	52.435
1108.01	18.925	24	5.613	0.2597	
1112.01	37.808	10	5.134	0.1537	
1113.01	25.935	16	4.896	0.2526	1.48
1113.02	83.441	6	2.871	0.1906	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
1115.01	12.992	31	5.501	0.3062	
1117.01	11.089	40	5.062	0.3357	
1145.01	30.585	16	9.449	0.1592	
1152.01	4.722	86	3.786	0.4089	
1159.01	64.617	6	3.969	0.0786	
1160.01	13.214	33	5.447	0.3057	
1165.01	7.054	61	4.231	0.3902	
1175.01	31.596	16	4.256	0.2801	
1176.01	1.974	200	5.303	0.4106	
1177.01	3.306	120	7.162	0.3734	
1187.01	0.371	1160	5.402	0.4273	
1191.01	8.173	19	4.732	0.2758	
1194.01	8.708	15	5.682	0.1899	
1198.01	16.089	28	3.486	0.3696	7.272
1198.03	35.678	12	4.698	0.2053	
1199.01	53.528	8	7.045	0.0549	
1203.01	31.882	9	3.878	0.116	0.959
1203.03	48.656	8	4.263	0.1678	
1205.01	8.639	37	3.912	0.3759	
1207.01	13.735	33	4.523	0.3641	
1210.01	14.554	26	4.115	0.3406	
1215.01	17.324	22	6.215	0.2291	5.847
1215.02	33.007	12	6.976	0.1325	
1216.01	11.131	42	5.533	0.3294	
1218.01	29.619	14	3.837	0.2793	
1221.01	30.157	14	5.809	0.2662	3.767
1221.02	51.081	9	3.814	0.2039	
1227.01	2.155	187	6.104	0.3993	
1236.01	35.746	13	5.468	0.1773	
1236.02	12.31	34	4.721	0.3376	3.08
1238.01	27.072	15	4.108	0.2721	
1241.01	21.406	21	4.669	0.2851	
1241.02	10.501	43	8.683	0.2288	10.955
1245.01	13.72	31	3.782	0.3595	
1246.01	19.037	23	6.258	0.2474	
1257.01	86.648	6	8.153	0.0192	
1258.01	36.338	13	4.713	0.2223	
1258.02	14.646	28	4.105	0.3426	2.7
1264.01	14.132	34	3.556	0.3749	
1266.01	11.419	38	4.268	0.3574	
1270.01	5.729	74	5.271	0.3786	21.274

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
1270.02	11.608	40	11.184	0.1615	
1273.01	40.058	11	5.239	0.1638	
1275.01	50.285	9	3.723	0.2066	
1276.01	22.789	22	4.427	0.3092	
1278.01	24.806	17	5.995	0.1934	5.405
1278.02	44.346	11	3.196	0.2982	
1279.01	14.374	32	3.078	0.3858	
1281.01	49.477	10	4.859	0.1586	
1282.01	30.864	14	4.818	0.2211	
1283.01	8.092	55	3.666	0.3937	
1285.01	0.937	453	11.112	0.3897	
1298.01	11.008	39	4.052	0.3713	
1299.01	52.5	7	5.293	0.0874	
1301.02	37.514	9	6.412	0.0577	
1302.01	55.638	8	4.087	0.1501	
1303.01	34.296	14	3.987	0.2766	
1307.01	44.851	10	5.287	0.1574	
1307.02	20.343	22	3.945	0.3219	1.296
1308.01	23.585	15	4.952	0.2377	
1309.01	10.117	44	4.615	0.3602	
1310.01	19.13	21	5.573	0.3295	
1311.01	83.576	6	5.116	0.0496	
1314.01	8.575	41	4.94	0.3452	
1315.01	6.846	64	4.406	0.3815	
1320.01	10.507	15	3.552	0.2752	
1323.01	3.99	41	5.003	0.3416	
1325.01	10.035	37	5.234	0.3289	
1329.01	33.199	12	5.657	0.1675	
1332.01	19.306	13	3.248	0.2899	
1336.02	15.573	23	5.858	0.246	
1353.02	34.543	13	3.598	0.2544	
1355.01	51.929	9	5.988	0.2621	
1357.01	3.014	82	4.682	0.3899	
1358.01	5.645	43	4.735	0.355	8.864
1358.02	8.744	28	4.402	0.3326	
1360.01	36.77	12	3.045	0.2988	
1360.02	14.59	30	5.709	0.2903	3.984
1361.01	59.879	8	5.967	0.0669	
1362.01	20.513	12	5.73	0.1639	
1364.01	20.833	17	5.057	0.2442	
1366.01	19.254	25	4.485	0.3129	7.987

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
1366.02	54.157	7	8.423	0.0381	
1372.01	69.668	6	3.87	0.0447	
1376.01	7.139	56	5.662	0.3533	
1378.01	19.302	22	4.456	0.3092	
1382.01	4.202	84	10.693	0.2658	
1385.01	18.61	23	4.154	0.3364	
1387.01	23.8	20	4.582	0.2544	
1391.01	7.981	54	4.578	0.3743	
1393.01	1.695	144	4.815	0.4061	
1397.01	6.247	37	4.128	0.3614	
1398.01	5.823	42	4.626	0.3604	
1399.01	8.75	28	5.261	0.2941	
1403.01	18.754	25	3.731	0.351	
1406.01	11.361	38	3.158	0.3867	
1408.01	14.534	32	5.05	0.3293	
1409.01	16.56	25	3.655	0.3516	
1410.01	15.75	28	3.523	0.3609	
1412.01	37.813	12	3.337	0.2949	
1413.01	12.645	34	4.003	0.3607	
1419.01	1.336	259	4.771	0.4178	
1420.01	13.352	17	4.848	0.2381	
1422.01	5.842	61	4.296	0.3867	2.365
1422.02	19.85	21	5.35	0.2671	
1426.01	38.873	9	5.537	0.1364	
1430.01	10.475	41	4.617	0.3567	2.319
1430.02	22.93	18	5.732	0.2174	0.995
1430.03	77.482	6	3.702	0.1348	
1433.01	19.808	19	4.692	0.2828	
1435.01	40.716	11	5.543	0.1583	
1435.02	10.446	44	5.548	0.3396	0.91
1436.02	13.751	28	8.933	0.1815	
1438.01	6.911	67	7.013	0.3336	
1444.01	44.932	9	4.114	0.2116	
1445.01	7.169	49	6.721	0.3236	
1448.01	2.487	167	5.286	0.4056	
1452.01	1.152	348	7.463	0.4069	
1456.01	7.887	33	4.989	0.3246	
1457.01	8.028	32	4.029	0.3503	
1458.01	8.981	23	5.567	0.2516	
1459.01	0.692	610	4.835	0.4255	
1465.01	9.771	40	4.397	0.3583	

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
1470.01	15.434	18	4.405	0.2835	
1472.01	85.351	6	2.555	0.2397	
1473.01	23.02	11	6.224	0.1389	
1475.02	9.513	29	5.699	0.3054	
1476.01	56.362	6	7.761	0.0599	
1480.01	20.381	22	3.186	0.361	
1481.01	5.101	50	5.347	0.35	
1486.02	30.184	15	3.884	0.2661	
1489.01	16.005	30	5.568	0.2906	
1495.01	15.595	27	3.648	0.3482	
1499.01	14.164	32	5.423	0.3165	
1506.01	40.43	11	4.811	0.1765	
1507.01	21.36	19	3.83	0.3056	
1508.01	22.047	20	4.376	0.3018	
1512.01	9.042	42	5.111	0.3347	
1516.01	20.554	21	3.893	0.3256	
1517.01	40.068	9	3.286	0.2464	
1518.01	27.506	17	5.779	0.2097	
1520.01	18.458	25	4.165	0.3283	
1521.01	25.941	14	3.252	0.2637	
1522.01	33.386	14	4.023	0.2744	
1525.01	7.714	50	3.737	0.388	
1529.01	17.976	23	10.896	0.1077	
1530.01	12.985	25	3.918	0.3485	
1532.01	18.115	22	4.094	0.3207	
1534.01	20.422	22	3.648	0.3494	
1540.01	1.208	340	8.227	0.4031	
1541.01	2.379	149	6.064	0.3958	
1543.01	3.964	112	8.655	0.3451	
1546.01	0.918	438	10.747	0.3912	
1547.01	30.694	6	11.292	0.0024	
1549.01	29.481	16	4.831	0.2489	
1553.01	52.759	9	7.513	0.0597	
1557.01	3.296	106	5.185	0.3958	4.8
1557.02	9.653	38	4.91	0.3405	
1557.03	5.316	66	4.706	0.3811	2.261
1560.01	31.569	14	5.218	0.2101	
1561.01	9.086	26	5.659	0.2685	
1563.01	5.487	42	3.348	0.3871	4.173
1563.02	8.291	27	3.976	0.3472	3.514
1563.04	16.74	15	5.062	0.225	

Acknowledgements

Funding for this mission is provided by NASA’s Science Mission Directorate. J.H.S acknowledges support by NASA under grant NNX08AR04G issued through the Kepler Participating Scientist Program. D. C. F. and J. A. C. acknowledge support for this work was provided by NASA through Hubble Fellowship grants #HF-51272.01-A and #HF-51267.01-A awarded by the Space Telescope Science Institute, operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555.

REFERENCES

- E. Agol, J. Steffen, R. Sari, and W. Clarkson. *MNRAS*, 359:567–579, May 2005.
- S. Ballard et al. *ApJ*, 743:200, December 2011.
- N.M. Batalha et al. *ApJS submitted*, 2012.
- W. J. Borucki et al. *ApJ*, 736:19–+, July 2011.
- J. A. Carter, E. Agol, et al. *Science*, In press, 2012.
- W. D. Cochran et al. *ApJS*, 197:7, November 2011.
- D. C. Fabrycky, E. B. Ford, J.-H. Steffen, et al. *ApJ*, 750, 114, 2012.
- E. B. Ford, D. C. Fabrycky, J.-H. Steffen, et al. *ApJ*, 750, 113, 2012a.
- E. B. Ford et al. *ApJS*, 197:2–+, November 2011.
- E.B. Ford et al. *ApJ submitted*, 2012b.
- J. S. Heyl and B. J. Gladman. *MNRAS*, 377:1511–1519, June 2007. doi: 10.1111/j.1365-2966.2007.11697.x.
- M. J. Holman and N. W. Murray. *Science*, 307: 1288–1291, February 2005.
- M. J. Holman et al. *Science*, 330:51–, October 2010.
- J. J. Lissauer et al. *Nature*, 470:53, January 2011.
- J. Miralda-Escudé. *ApJ*, 564:1019–1023, January 2002. doi: 10.1086/324279.
- J.F. Rowe et al. *ApJS to be submitted*, 2012.
- J. H. Steffen. *Ph.D. Dissertation*, University of Washington, 2006. arXiv:astro-ph/0609492.
- J. H. Steffen and E. Agol. *MNRAS*, 364:L96–L100, November 2005. doi: 10.1111/j.1745-3933.2005.00113.x.
- J.H. Steffen, E. B. Ford, D. C. Fabrycky, et al. *MNRAS*, 421, 2342, 2012a.
- J.H. Steffen et al. *PNAS*, 109, 21, 7982-7987, 2012b.
- G. M. Szabó, A. Pál, A. Derekas, A. E. Simon, T. Szalai, and L. L. Kiss. *ArXiv e-prints:1110.4231*, October 2011.
- B. Tingley et al. *A&A*, 536:L9, December 2011.

TABLE 1—*Continued*

KOI	Period (days)	N_t	SNR	p-value	$\Xi_{\max}^b$
1564.01	53.449	8	3.149	0.2498	7.314
1567.01	7.24	34	4.381	0.3468	
1567.03	17.327	15	5.093	0.254	
1569.01	13.752	32	3.884	0.3599	
1570.01	6.339	37	5.037	0.348	
1572.01	9.803	24	4.678	0.3003	2.277
1573.01	24.809	17	42.306	0.0079	
1576.01	10.416	41	5.558	0.3252	
1576.02	13.084	33	4.961	0.3293	
1581.01	29.547	14	11.533	0.0741	
1585.01	19.179	23	4.659	0.288	5.147
1587.01	52.972	8	13.883	0.0048	
1588.01	3.517	127	5.206	0.3998	
1589.01	8.726	52	5.428	0.3547	
1589.02	12.882	35	4.272	0.3619	
1589.03	27.435	17	3.362	0.326	3.005
1589.05	44.549	11	4.248	0.2159	
1591.01	19.657	24	3.765	0.334	
1595.01	40.109	12	3.693	0.2355	
1597.01	7.797	53	4.516	0.3695	
1598.01	56.475	6	3.353	0.1329	0.085
1599.01	20.419	21	11.668	0.085	
1601.01	10.351	37	3.666	0.3772	
1608.01	9.176	45	3.984	0.3747	
1609.01	41.698	10	4.577	0.2153	

^aThis table is published in its entirety in the electronic edition.

^b $\Xi_{\max}$ values are given for each KOI with its nearest exterior neighbor that satisfies the selection criteria for analysis (i.e., its exterior neighbor that is also reported in this table).